

**Kh.I.Strus,
A.M.Yashchenko**

Danylo Halytsky Lviv
National Medical Uni-
versity

Key words: hypothy-
roidism, skin, hair, se-
baceous gland, lectin
SNA, lectin WGA.

Received: 14.10.2012

Accepted: 24.11.2012

UDC 611.773/.78-018:547.96]:616.441-008.64.]-019-08

SIALOGLYCANS OF SKIN DERIVATIVES OF RAT OFFSPRING ON THE 10TH POST- NATAL DAY OF DEVELOPMENT AT THE BACKGROUND OF EXPERIMENTAL MERCAZOLIL-INDUCED HYPOTHYROIDISM OF MATERNAL ORGANISM

The study was conducted as a part of research works “Lectin markers and cytoplasmic signal molecules in the process of cellular differentiation and proliferation” (state registration 0700U00106), “Searching for new lectin preparations from the raw materials of the Carpathian region and for the ways of their use in biology and medicine” (state registration 0107U001048).

Summary. With the use of sialo-specific lectins WGA and SNA, to investigate the role of sialoglycans in differentiation of structural components of skin derivatives in the offspring of control and hypothyrotic rats on the 10th postnatal day of development. Hypothyroidism was modeled by introduced with food mercazolil (5 mg/kg of body weight). Sliced skin taken from the back of offspring of control and hypothyrotic females were fixed in 4% neutral formalin and embedded in paraffin. Preparations 5-7 μm thick were stained with hematoxylin and eosin to study their general morphology. Lectin conjugates with horse-raddish peroxidase were used to identify carbohydrate determinants. Lectin receptors were visualized in the system 3,3'-diaminobenzidin-H₂O₂. Reduced activity of metabolic processes at the background of thyroid hypofunction in the maternal organism was noticed to be associated with reduced morphogenesis as well as slight decrease of the expression of sialoglycans that act as signal molecules and influencing growth processes and differentiation of structural components of skin derivatives.

Citation: Strus KhI, Yashchenko AM. [Sialoglycans of skin derivatives of rat offspring on the 10th postnatal day of development at the background of experimental mercazolil-induced hypothyroidism of maternal organism]. *Morphologia*. 2012;6(4):83-9. Ukrainian.

© Strus Kh.I., Yashchenko A.M., 2012





Fig. 1. Thyroid gland. Hematoxylin & Eosin stainings. A – group of control. $\times 150$. Б – experimental group (after mercasolil administration). $\times 150$.




Fig. 2. Rat skin on the 10th day of postnatal development. Hematoxylin & Eosin stainings. A – hair follicles of the control group rats' offsprings. ×300. Б – hair follicles of the hypothyroid female rats' offsprings. ×300. B – hair follicle and sebaceous gland of the control group rats' offsprings. ×600. Г – hair follicle and sebaceous gland of the experimental group rats' offsprings. ×600.




Fig. 3. Expression of the SNA lectin receptors in the skin derivatives on the 10th day of postnatal development. A – hair follicles of the control group rats' offsprings. ×300. Б – hair follicles of the hypothyroid female rats' offsprings. ×300. B – SNA lectin receptors in the external epithelial vagina and the medulla of the hair root of the control group rats' offsprings. ×600. Г – decrease of the SNA lectin receptor expression in the medulla of the hair root in the hypothyroid female rats' offsprings. ×600. Д – binding the SNA lectin receptors to the surface of the sebocytes in the sebaceous glands of the control group rats' offsprings. ×600. E – slight decrease of the SNA lectin receptor expression on the surface of the sebocytes in the sebaceous glands in the hypothyroid female rats' offsprings. ×600.




Fig. 4. Expression of the WGA lectin receptors in the skin derivatives on the 10th day of postnatal development. A – expression of the WGA lectin receptors in the medulla of the hair root of the control group rats' offsprings. ×600. Б – slight expression of the WGA lectin receptors in the cuticle of the hair root of experimental animals. ×600. B – the WGA lectin receptors on the surface of the sebocytes and in the basal membrane of the terminal secretory portions of the sebaceous glands of the control group rats' offsprings. ×600. Г – decrease of the binding intensity in the sebocytes of the sebaceous glands of the experimental group of animals. ×600.

References:

Antonyuk VO. Lectyny ta yikh syrovynni dzherela [Lectins and the sources of raw materials]. Lviv: Kwart; 2005. 554 p. Ukrainian.

Bernik NV, Oliinyk IYu, Lavriv LP. [Human morphology and lectinohistochemistry]. Clin and experim pathol. 2010;9(3):138-43. Ukrainian.

Voloshin NA, Paschenko SN. [Role of lectins in the diagnosis and treatment of malignant tumors]. Zaporozhye Medical Journal. 2004;(3):93-6. Russian.

Lutsik AD, Detyuk YeS, Lutsik MD. Lektyny v gistokhimii [Lectins in histochemistry]. Lviv: Vyscha shkola; 1989. 144 p.

Lutsik OD, Benkston PV. [Heterogeneity of certain cell populations in rat detected by lectin histochemistry]. Acta Medica Leopoliensia. 1997;3(1-2):70-9. Ukrainian.

Lutsyk SA, Yashchenko AM. [Experimental hypothyroidism induce increased exposure of alpha-LFuc, beta-DGal and DGalNAc carbohydrate determinants in rat adrenal glands]. World of medicine and biology. 2012;8(2):123-7. Ukrainian.

Preima ChI, Yashchenko AM. [Role of glycoconjugates in the processes of skin morphogenesis in the offspring with underlying maternal organism's hypothyroidism]. *Buk Med Herald*. 2009;13(4):232-6. Ukrainian.

Tabachnyuk NV, Oliinyk IYu, Lavriv LP. [Lectinohistochemical studies and embryogenesis]. *Klinichna anatomiya ta operatyvna khirurgiya*. 2010;9(3):95-100. Ukrainian.

Brooks SA, Harris A, editors. *Breast Cancer Research Protocols (Methods in Molecular Medicine)*. NJ: Humana Press; 2006. 517 p.

Forslind B, Albano R. *Skin, hair, and nails*. CRC Press; 2003. 469 p.

Ishii M, Tsukise A, Meyer W. Lectin histochemistry of glycoconjugates in the feline hair follicle and hair. *Ann Anat*. 2001 Sep;183(5):449-58. Cited in: PubMed; PMID: 11677811.

Krinke G. *The laboratory rat*. London: Academic Press; 2000. 756 p.

Roth J. Lectins for histochemical demonstration of glycans. *Histochem Cell Biol*. 2011 Aug;136(2):117-30. doi: 10.1007/s00418-011-0848-5. Epub 2011 Jul 31. Cited in: PubMed; PMID: 21805335.

Sharon N. Lectins: carbohydrate-specific reagents and biological recognition molecules. *J Biol Chem*. 2007 Feb 2;282(5):2753-64. Cited in: PubMed; PMID: 17145746.

Tsujio M, Yoshioka K, Satoh M, Watahiki Y, Mutoh K. Skin morphology of thyroidectomized rats. *Vet Pathol*. 2008 Jul;45(4):505-11. doi: 10.1354/vp.45-4-505. Cited in: PubMed; PMID: 18587098.

Varki A. Glycan-based interactions involving vertebrate sialic-acid-recognizing proteins. *Nature*. 2007;446(7139):1023-9.