

**M.A.Ovdii**

Bogomolets National  
Medical University,  
Kyiv, Ukraine

**Key words:** morpho-  
functional parameters,  
overweight, young  
women.

*Received: 07.11.2013*  
*Accepted: 23.12.2013*

UDC:616-056.52-055.2-053.81[616-008+616-071.3]

## **DYNAMICS OF MORPHOFUNCTIONAL PARAMETERS YOUNG WOMEN IN THE PROCESS OF CORRECTION OVER- WEIGHT**

*The study was performed as a part of the research work “Physical preparedness and quality of life of young people depending on body weight” (state registration number 011U001410).*

**ABSTRACT. Background.** Currently, much attention is paid to the problem of correction and prevention overweight and obesity in young people using different methods of physical activity, however, the use of the walking with breathing exercises presented in the literature is not enough. **Objective.** The present study investigated the dynamic of morphological and functional indicators in young women with overweight under the influence of the walking with breathing exercises. **Methods.** 23 young women (21-25 years) participated in the present study and performed walking with breathing exercises four times per week for 12 weeks. Before and after the intervention, measurements of weight, body mass index, waist, abdomen and hips circumferences, body composition, and functional parameters. **Results.** After 12 weeks found a positive trend in anthropometric and functional parameters, reduced body weight by 8,7% ( $p < 0,01$ ), body fat by 6,9% ( $p < 0,01$ ), waist and abdomen circumferences by 8,2 % and 6,9% ( $p < 0,01$   $p < 0,05$  respectively), increased lung capacity by 15,7% ( $< 0,01$ ), adaptation of the cardiovascular system to the conditions of hypoxia and other functional parameters. **Conclusion.** The present study suggests that walking with breathing exercises improves the morphofunctional parameters overweight young women.

© M.A.Ovdii, 2013  
✉ [omarija1@rambler.ru](mailto:omarija1@rambler.ru)

### **Citation:**

Ovdii M.A. [Dynamics of morphofunctional parameters young women in the process of correction overweight]. *Morphologia*. 2013;7(4):58-61. Russian.

## *References:*

1. Ametov AS. [Obesity epidemic of the XXI century]. Therapeutic Archives. 2008;74(10):5-7. Russian.
2. Brock DW, Thomas O, Cowan CD, Allison DB, Gaesser GA, Hunter GR. Association between insufficiently physically active and the prevalence of obesity in the United States. J Phys Act Health. 2009;6(1) :1-5. PMID:19211952; PMCID: PMC2659323
3. Allison DB, Newcomer JW, Dunn AL, Blumenthal JA, Fabricatore AN, Daumit GL. Obesity among those with mental disorders: a National Institute of Mental Health meeting report. Am J Prev Med. 2009;36(4):341-50. PMID:19285199
4. Ogden CL, Yanovski SZ, Carroll MD, Flegal KM. The epidemiology of obesity. Gastroenterology. 2007;132(6):2087-102. PMID: 17498505
5. Edwards AM, Walker RE. Inspiratory muscle training and endurance: a central metabolic control perspective. Int J Sports Physiol Perform. 2009;4(1):122–128. PMID: 19417233
6. Thompson D, Karpe F, Lafontan M, Frayn K. Physical activity and exercise in the regulation of human adipose tissue physiologi. Physiol Rev. 2012;92(1):157-91. PMID: 22298655
7. Ball K, Brown W, Crawford D. Who does not gain weight? Prevalence and predictors of weight maintenance in young women. Int J Obes Relat Metab Disord. 2002;26(12):1570-8. PMID: 12461673
8. Wuyam B, Verg`es S. Respiratory muscle endurance training in obese patients. Int J Obes (Lond). 2011;35(5):692-9. PMID: 20805830