

I.G.Savka

Bukovinian State
Medical University;
Chernivtsi, Ukraine

Key words: structural and functional features, long bones, lower extremity fractures, morphological features.

Received: 26.11.2013
Accepted: 20.12.2013

UDC 616.718-001.5-091:611.718

SUBSTANTIATION OF THE INFLUENCE OF STRUCTURAL-FUNCTIONAL PECULIARITIES OF CERATIN PORTIONS OF TUBULAR BONES OF THE LOWER EXTREMITY UPON THE FORMATION OF MORPHOLOGICAL SIGNS OF FRACTURES

The study was performed as a part of research work "Patterns of perinatal anatomy and embryotopography. Determination of sex and age peculiarities of structure and topographic anatomical interrelations of organs and structures in the human ontogeny" (state registration number 0110U003078).

ABSTRACT. Background. Forensic doctors and traumatologists deal with more and more cases of atypical fractures, minimal injuries of unknown etiology and mechanism. The relationship between external and internal factors in these cases still remains understudied. **Objective.** To study the influence of structural-functional peculiarities of certain portions of tubular bones of the lower extremities upon the formation of morphological signs of fractures which might ensure an objective retrospective detection of the mechanisms of their formation. **Methods.** The objects of our studies were expert observations of 128 cases with injuries of the femoral bone, tibia and fibula. The obtained results were statistically processed with the use of single-factor dispersion analysis, primary analysis with Kettel's test and multi-factor analysis. **Results.** Various portions of the long tubular bones of the lower extremity were found to have a number of structural-functional peculiarities. Detection of morphological signs of long tubular bones fractures enables to identify the character of the injury and find the mechanism of its occurrence. Depending on the bone portion the action of mechanical force of the same nature leads to fractures with different morphological characteristics. The value of emerging angle of wedge-shaped fissures is of a diagnostic importance among macro-architectural parameters. **Conclusion.** Perspective is further in-depth study of the relationships between the main structural components of bone formation and the patterns of morphological characteristics formation in fractures of various bones of the human skeleton.

© I.G.Savka, 2013

Citation:

Savka IG. [Substantiation of the influence of structural-functional peculiarities of ceratin portions of tubular bones of the lower extremity upon the formation of morphological signs of fractures]. *Morphologia*. 2013;7(4):62-6. Ukrainian.

References:

1. Leonov SV, Leonova EN. [Specific features of the injuries to diaphyses of long tubular bones in case of skew bending]. *Forensic Medical Examination*. 2011; 54(6): 13-5. Russian.
2. Nagornov MN, Osipenkova-Vichtomova TK. [Forensic medical aspects of bone tissue injuries and pathology]. *Forensic Medical Examination*. 2012; 55(1): 41-4. Russian.
3. Pigolkin YuI, Nagornov MN. [Forensic evaluation of bone fractures]. *Forensic Medical Examination*. 2005; 48(6): 39-42. Russian.
4. Yankovsky VE. [The role of stretching in the process of fractures formation]. *Forensic Medical Examination*. 2008; 51(2): 3-6. Russian.
5. Keaveny TM, Bouxsein ML. Theoretical implications of the biomechanical fracture threshold. *J Bone Miner Res*. 2008 Oct;23(10):1541-7. doi: 10.1359/jbmr.080406. Cited in: PubMed; PMID: 18410232; PMCID: PMC2684155.
6. Kosmopoulos V, Keller TS. Predicting trabecular bone microdamage initiation and accumulation using a non-linear perfect damage model. *Med Eng Phys*. 2008 Jul;30(6):725-32. Epub 2007 Sep 18. Cited in: PubMed; PMID: 17881275.
7. Nagaraja S, Couse TL, Guldborg RE. Trabecular bone microdamage and microstructural stresses under uniaxial compression. *J Biomech*. 2005 Apr;38(4):707-16. Cited in: PubMed; PMID: 15713291.
8. Sroga GE, Karim L, Colón W, Vashishth D. Biochemical characterization of major bone-matrix proteins using nanoscale-size bone samples and proteomics methodology. *Mol Cell Proteomics*. Sep 2011; 10(9): M110.006718. doi: 10.1074/mcp.M110.006718. Cited in: PubMed; PMCID: PMC3186195.
9. Waldorff EI, Goldstein SA, McCreadie BR. Age-dependent microdamage removal following mechanically induced microdamage in trabecular bone in vivo. *Bone*. 2007 Feb;40(2):425-32. Epub 2006 Oct 19. Cited in: PubMed; PMID: 17055351.
10. Wang CJ, Liu HC, Fu TH. The effects of extracorporeal shockwave on acute high-energy long bone fractures of the lower extremity. *Arch Orthop Trauma Surg*. 2007 Feb;127(2):137-42. Epub 2006 Oct 13. Cited in: PubMed; PMID: 17053946.
11. Yu XZ, Guo YM, Li J, Zhang YQ, He RX. Finite element analysis of impact loads on the femur. *Chin J Traumatol*. 2007 Feb;10(1):44-8. Cited in: PubMed; PMID: 17229350.