

**P.I.Gritsenko
N.S.Petruk
A.V.Samoylenko
I.V.Tverdokhle**

State establishment
“Dnipropetrovsk
Medical Academy of
Health Ministry of
Ukraine”

Key words: primary chronic periodontitis, secondary chronic periodontitis, histological structure, ultrastructure.

*Received: 14.05.2014
Accepted: 03.06.2014*

UDC 616.314.17-036.1-008.811.9-08:615.37

CELLULAR COMPOSITION AND ULTRASTRUCTURE OF PERIAPICAL GRANULATION TISSUE IN PRIMARY AND SECONDARY CHRONIC PERIODONTITIS

The study was performed as the part of research work “Development and introduction of new methods for diagnosis, treatment and prevention of dental caries and its complications, diseases of periodontium and oral mucosa” (state registration number 0110U003018).

ABSTRACT. Background. There are no complete data on the occurrence of bacteria in the outbreak of chronic inflammation a consequence of their high virulence, or the result of a defect of local mechanisms of immune protection. To answer these questions, as well as to evaluate the nature, severity changes periapical tissues of the tooth and the activity of the inflammatory process in the apex of the tooth root is possible only during the morphological study of biopsy material. **Objective.** The aim of this work was to study the cellular composition and ultrastructural features of the granulation tissue formed during the current the primary and secondary apical periodontitis. **Methods.** There were included in the histological examination of the teeth 78, who underwent resection of the apex, followed by a fence of histological material. For ultrastructural studies of granuloma periodontal tissue samples were fixed with 2.5% glutaraldehyde solution, material embedded in Epon-812 («SPI-Pon™ 812 Epoxy Embedding Kit», USA). The study was conducted using a transmission electron microscopy. **Results.** In the development of destructive periodontal inflammation are actively involved both specific and non-specific mechanisms of immunity. **Conclusion.** In primary periodontitis granulation tissue contains a moderate amount of white blood cells of various types, as well as single macrophages and plasma cells with reduced functional activity. In secondary periodontitis granuloma contains numerous mononuclear macrophages, plasma cells and giant multinucleated cells with increased activity, which corresponds to the morphological picture of the formation of immune granulomas.

© P.I.Gritsenko, N.S.Petruk, A.V.Samoylenko, I.V.Tverdokhle, 2014
✉ gritsenko-pavlo@meta.ua

Citation:

Gritsenko PI, Petruk NS, Samoylenko AV, Tverdokhle IV. [Cellular composition and ultrastructure of periapical granulation tissue in primary and secondary chronic periodontitis]. *Morphologia*. 2014;8(2):14-9. Russian.




Fig.1. Plot of the plasma cell (ПК) within the periapical granuloma in patients with primary chronic granulomatous periodontitis. Low activity of the nucleus, insignificant osmiophily of "light court" (arrow). Moderate synthetic activity of granular reticulum. x8000.




Fig.2. Lysosomes (arrow) in the cytoplasm of macrophage (МФ) bordering the collagen fibers (КВ) of granulation tissue in patient with primary chronic granulomatous periodontitis. x8000.




Fig.3. Giant multinucleated cell in the composition of periapical granuloma in patient with secondary chronic granulomatous periodontitis. $\times 4000$.

References:

1. Walton RE. Routine histopathologic examination of endodontic periradicular surgical specimens – is it warranted? *Oral Surg Endod.* 1998;86(5):505.
2. Haapasalo H, Siren E, Waltimo T, Orstavik D, Haapasalo M. Inactivation of local root canal medicaments by dentine: an in vitro study. *Int End J.* 2000;2(33):126.
3. Nair PNR. Pathogenesis of apical periodontitis and the causes of endodontic failures. *Crit Rev Oral Biol Med.* 2004;15:348-81.
4. Rabuhina HA, Grigor'yan AC, Grigor'yants LA, Badaljan VA [Comparison of radiographic, clinical and morphological parameters with radicular destructive lesions]. *Clinical Dentistry.* 1999;3:24-7. Russian.
5. Tanomaru-Filho M, Poliseli-Neto A, Leonardo MR, Silva LAB, Tanomaru JMG, Ito IY. Methods of experimental induction of periapical inflammation. Microbiological and radiographic evaluation. *Int Endod J.* 2005;38:477-82.
6. Kabak SL, Kabak YuS, Anishchenko SL, Kolb YeL. [Cellular composition of periapical granuloma]. *Vestsi NAS Belarusi. Ser med navuk.* 2009;1:44-50. Russian.

7. Sarkisov DS, Perov YL. Mikroskopicheskaya tekhnika. Rukovodstvo dlya vrachey i laborantov [Microscopic technique. A guide for doctors and laboratory technicians]. Moscow: Medcine; 1996. 544 p. Russian.

8. Weekly B. Elektronnaya mikroskopiya dlya nachinayushchikh [Electron microscopy for beginners]. Moscow: Mir; 1975. 324 p. Russian.

9. Hernandez-Pando R, Bornstein QL, Aguilar Leon D, Orozco EH, Madrigal VK, Martinez CE. Inflammatory cytokine production by immunological and foreign body multinucleated giant cells. *Immunology*. 2000;100(3):352-8.

10. Anderson JM. Multinucleated giant cells. *Curr Opin Hematol*. 2000;7:40-7.

11. Zhu XW, Friedland JS. Multinucleate giant cells and the control of chemokine secretion in response to Mycobacterium tuberculosis. *Clin Immunol*. 2006;120:10-20.