

**A.O.Makarchuk**

Zaporizhzhia State  
Medical University

**Key words:** skin,  
involutive changes,  
malasseziasis,  
ultrastructure.

*Received: 24.02.2015*  
*Accepted: 23.03.2015*

UDC 616.5:612.66]-06:616022]-085

## **THE DYNAMICS OF MORPHOLOGICAL CHANGES OF FACIAL SKIN IN PATIENTS WITH MALASSEZIASIS AT STAGES OF ANTIMYCOTIC AND ANTI-AGING THERAPY**

**ABSTRACT. Background.** Many questions about the dependence of pathological changes of the skin to age-appropriate background malassezia destruction require clarification and further advance, as well as the dependence of the effectiveness of anti-aging interventions of antimycotic therapy. **Objective.** The purpose of this study was to determine the morphological changes of the skin in patients with malasseziasis on stages of antimycotic and anti-aging therapy. **Methods.** In 90 patients with involutive changes in facial skin, including 60 patients with malasseziasis, using electron microscopy carried determine the state of the skin. **Results.** Found that specific antifungal therapy for 1 month prior to surgical intervention in the face provides the normalization of skin structures damaged by keratomycosis, through emulation of mitotic activity of keratinocytes in germinal zone and optimizing epidermal differentiation restricted vacuolization of epithelial cells. Prolongation of local and systemic antimycotic therapy within 1 month after the intervention was accompanied by reduction of involutive morphological changes of the skin and prevents development of malasseziasis recurrent for 1 year. **Conclusion.** Specific antimycotic therapy in patients with malasseziasis is essential for achieving high performance anti-aging activities.

© **A.O.Makarchuk, 2015**  
✉ **makarchuk.art@gmail.com**

### **Citation:**

Makarchuk AO. [The dynamics of morphological changes of facial skin in patients with malasseziasis at stages of antimycotic and anti-aging therapy]. *Morphologia*. 2015;9(1):39-46. Ukrainian.




Fig. 1. Dynamics of changes in the numerical density of keratinocyte nuclei in germinative layers of epidermis ( $10^3 \times \text{mm}^{-2}$ ) in patients aged 33-40 years. Note: Asterisk (\*) indicates the reliable difference between the main clinical group and the comparative group.




Fig. 2. Dynamics of changes in the numerical density of keratinocyte nuclei in germinative layers of epidermis ( $10^3 \times \text{mm}^{-2}$ ) in patients aged 41-50 years. Note: Asterisk (\*) indicates the reliable difference between the main clinical group and the comparative group.




Fig. 3. Dynamics of changes in the numerical density of keratinocyte nuclei in germinative layers of epidermis ( $10^3 \times \text{mm}^{-2}$ ) in patients aged 51-57 years. Note: Asterisk (\*) indicates the reliable difference between the main clinical group and the comparative group.




Fig. 4. Dynamics of changes in the general thickness of basal, spinous and granular layers of epidermis ( $\mu\text{m}$ ) in patients aged 33-40 years.




Fig. 5. Dynamics of changes in the general thickness of basal, spinous and granular layers of epidermis ( $\mu\text{m}$ ) in patients aged 41-50 years. Note: Asterisk (\*) indicates the reliable difference between the main clinical group and the comparative group.




Fig. 6. Dynamics of changes in the general thickness of basal, spinous and granular layers of epidermis ( $\mu\text{m}$ ) in patients aged 51-57 years. Note: Asterisk (\*) indicates the reliable difference between the main clinical group and the comparative group. ( $p < 0,05$ ).




Fig. 7. Dynamics of changes in the keratinocyte mitotic index (%) in patients aged 33-40 years. Note: Asterisk (\*) indicates the reliable difference between the main clinical group and the comparative group.




Fig. 8. Dynamics of changes in the keratinocyte mitotic index (%) in patients aged 41-50 years. Note: Asterisk (\*) indicates the reliable difference between the main clinical group and the comparative group.




Fig. 9. Dynamics of changes in the keratinocyte mitotic index (%) in patients aged 51-57 years. Note: Asterisk (\*) indicates the reliable difference between the main clinical group and the comparative group.




Fig. 10. Dynamics of changes in the keratinocyte vacuolation grade (%) in the germinative zone of patients aged 33-40 years. Note: Asterisk (\*) indicates the reliable difference between the main clinical group and the comparative group.




Fig. 11. Dynamics of changes in the keratinocyte vacuolation grade (%) in the germinative zone of patients aged 41-50 years. Note: Asterisk (\*) indicates the reliable difference between the main clinical group and the comparative group.




Fig. 12. Dynamics of changes in the keratinocyte vacuolation grade (%) in the germinative zone of patients aged 51-57 years. Note: Asterisk (\*) indicates the reliable difference between the main clinical group and the comparative group.

## References

1. Jiang LI, Stephens TG, Goodman R. SWIRL, a clinically validated, objective, and quantitative method for facial wrinkle assessment. *Skin Res Technol.* 2013;19(4):492-8.
2. Pessa JE, Nguyen H, John GB, Scherer PE. The anatomical basis for wrinkles. *Aesthet Surg J.* 2014;34(2):227-34.
3. El-Domyati M, Medhat W, Abdel-Wahab HM. Forehead wrinkles: a histological and immunohistochemical evaluation. *J Cosmet Dermatol.* 2014;13(3):188-94.
4. Park JY, Jang YH, Kim YS. Ultrastructural changes in photorejuvenation induced by photodynamic therapy in a photoaged mouse model. *Eur J Dermatol.* 2013;23(4):471-7.
5. Ovchinnikov RS, Manoyan MG, Gaynullina AG. Fungi malassezia in animal diseases: clinical forms, diagnosis. *VetPharma.* 2013;3(14):18-22. Russian.
6. Sosa ML, Rojas F, Mangiaterra M, Giusiano G. Prevalence of Malassezia species associated with seborrheic dermatitis lesions in patients in Argentina. *Rev Iberoam Micol.* 2013;30(4):239-42.
7. Nikityuk BA, Chtetsov VP, editors. [Human morphology]. 2nd ed. Moscow: Izd-vo MGU; 1990. 368 p. Russian.
8. Mironov AA, Komissarchik YuYa, Mironov VA. [Electron microscopy methods in biology and medicine : Methodological Guide]. St Petersburg: Science; 1994. 400 p. Russian.
9. Kuo J. Electron microscopy: methods and protocols. Totowa, New Jersey: Humana Press Inc. 2007. 608 p.
10. Avtandilov GG. Meditsinskaya morfometriya [Medical morphometry]. Moscow: Meditsina; 1990. 384 p. Russian.
11. Méndez-Vilas A, Rigoglio NN, Mendes Silva MV. Current microscopy contributions to advances in science and technology. Badajoz: Formatex; 2012. 1523 p.
12. Lakin GF. Biometriya [Biometrics: 4th ed.]. Moscow : Vysshaya shkola; 1990. 352 p. Russian.