

O.M.Besedin¹
K.V.Tsygankov¹
I.V.Kuzhevsky¹
P.O.Gritsenko²

1 Dnipropetrovsk City
Clinical Hospital №4

2 State institution
“Dnipropetrovsk
medical academy of
the Ministry of Health
of Ukraine”

Key words: vacuum
therapy,
morphological
characteristics,
capillaries,
fibroblasts.

Received: 20.05.2015
Accepted: 23.06.2015

UDC 616.379-008.64:617.586-06-08:615.811:616-001.4:576.31

CELL TRANSFORMATION OF WOUNDS IN DIABETIC FOOT PATIENTS UNDER VACUUM THERAPY

ABSTRACT. Background. The presence of micro- and macrovascular complications of bacterial contamination of the formation of biofilms, reduce the number of growth factors, chronic wound healing, enzymatic imbalance in the wound significantly impede treatment of patients with purulent necrotic complications of diabetic foot syndrome. **Objective.** The aim of our study is to evaluate the morphological changes at the cellular level in wounds in patients with diabetic foot syndrome on the background of vacuum therapy compared to traditional treatment. **Methods.** Morphological analysis of treatment of 12 patients with purulent complications of diabetic foot syndrome with foot lesions stages II-III by Wagner classification was carried out. **Results.** The study is based on the results of treatment of patients with purulent-necrotic complications of diabetic foot syndrome in terms of the morphological changes at the cellular level in biopsies of wounds. On the background of vacuum therapy in patients with wounds have investigated the maximum value of the average number of capillaries in the view field was $6,80 \pm 0,34$. Positive changes in microhemodynamics wounds on the background vacuum explained qualitative characteristics of the capillaries. The maximum mean diameter of fibroblast nuclei in the field of patients who underwent vacuum wound therapy defects was $82,89 \pm 4,51$, in patients with conventional wound conducting the figure was $60,18 \pm 3,17$. The trend of improving morphological parameters investigated at the cellular level, explains the effectiveness of the reparation of the vacuum therapy of wounds. **Conclusion.** Vacuum therapy optimizes microhemodynamics in wounds permeability of the walls of blood vessels newly formed granulation tissue, prevents the formation of protein deposits in the vessel walls and in the intercellular matrix of granulation tissue. This facilitates intercellular communication and promotes normal path of regenerative processes.

© O.M.Besedin, K.V.Tsygankov, I.V.Kuzhevsky, P.O.Gritsenko,
2015
✉ bam-86@mail.ru

Citation:

Besedin OM, Tsygankov KV, Kuzhevsky IV, Gritsenko PO. [Cell transformation of wounds in diabetic foot patients under vacuum therapy]. *Morphologia*. 2015;9(2):18-23. Ukrainian.




Fig. 1. Morphological features of wound biopsy in patients after conducted treatment. A. Granulation tissue after standart treatment. Sleeve-like thickening of capillary walls, condensation of intercellular substance with tendency to hyalinosis. Mallory-Slinchenko staining. B. Granulation tissue after VAC-therapy. Obliteration of capillaries, cords of endothelium-like cells. × 400. Hematoxylin&Eosin staining. C - Granulation tissue after VAC-therapy. Bundles of collagen fibers “scattered” around the obliterated capillaries. Mallory-Slinchenko staining. ×400.

References

1. Besedin OM, Malyk YY, Karpenko LI, Sotnikov SV, Jwanski LV, Pundik GM [Vacuum instillation therapy in patients with diabetic foot syndrome]. *Clinical Surgery*. 2014;11(3):8-10. Ukrainian.
2. Zaytseva EL, Tokmakova AYu. [Effects of growth factors and cytokins on soft tissue regeneration in patients with diabetes mellitus]. *Diabetes Mellitus*. 2014;(1):57-62. Russian. doi: 10.14341/DM2014157-62.
3. Michailichenko PP Osnovi vaccum terapyy [Fundamentals of vacuum therapy]. St. Petersburg: Sova; 2005. 318 p. Russian.
4. Yaremkevych RV [Clinical and morphological characteristics and features of diagnosis septic lesions in diabetic foot syndrome]. *Ukrainian Medical Almanac*. 2006;9:23-7. Ukrainian.
5. Zaytseva EL, Doronina LP, Molchkov RV, Voronkova IA, Mitish VA, Tokmakova AYu. [Effect of negative pressure therapy on repair of soft tissues of the lower extremities in patients with neuropathic and neuroischaemic forms of diabetic foot syndrome]. *Diabetes Mellitus*. 2014;(3):113-121. Russian. doi: 10.14341/DM20143113-121.

6. Boulton A, Cavanagh P, Raymann G. New and alternative treatments for diabetic foot ulcers: hormones and growth factors. The foot in diabetes, 4th edition. New York (USA): John Willey& Sons, Ltd; 2006. 360 p.

7. Dinh T, Tecilazich F, Kafanas A, Doupis J, Gnardellis C, Leal E, Tellechea A, Pradhan L, Lyons TE, Giurini JM, Veves A. [Mechanisms involved in the development and healing of diabetic foot ulceration]. Diabetes. 2012;61(11):2937-47. doi: 10.2337/db12-0227. PMID: 22688339; PMCID: PMC3478547

8. Erba P, Ogawa R, Ackermann M, Adini A. Angiogenesis in wounds treated by microdeformational wound therapy. Ann Surg. 2011;253:402-9.

9. Morykwas M, Faller B, Pearce D. [Effects of varying levels of subatmospheric pressure on the rate of granulation tissue formation in experimental wounds in swine]. Ann Plastic Surg. 2001;47(5):547-51.

10. Bassetto F, Lancerotto L, Salmaso R. Histological evolution of chronic wounds under negative pressure therapy. J Plast Reconstr Aesthet Surg. 2012;65(1):91-9.

11. Zaitsev VM, Lifyandsky VG, Marinkin VI [Applied Medical Statistics]. Moscow: Publishing House of the FOLIO; 2003. 432 p. Russian.