

V.Iu.Kravtsov ^{1,2,3}
N.V.Ibragimova ³
S.N.Nikonovich ⁴
E.A.Nadyrov ⁴
A.V.Rozhko ⁴

¹ Pushkin Leningrad State University

² The All-Russian state research institute of genetics and breeding of farm animals of the Russian Academy of Agricultural Sciences

³ The Federal State Institute of Public Health 'The Nikiforov Russian-Center of Emergency and Radiation Medicine', The Ministry of Russian Federation for Civil Defence, Emergencies and Elimination of Consequences of Natural Disasters

⁴ Republican Scientific and Practical Center for Radiation Medicine and Human Ecology, Republic of Belarus, Gomel

Key words:

thyroid cancer, nuclear abnormalities, chromosome bridges, Chernobyl accident.

Received: 22.10.2015

Accepted: 30.11.2015

UDC [616.444.-052.2 : 614.876] : 613.9 (476.2)

INTERNUCLEAR CHROMOSOME BRIDGES IN THYROCYTES OF PAPILLARY THYROID CANCER IN PATIENTS, SUBJECTED TO RADIOACTIVE IODINE ISOTOPES DURING FIRST MONTHS AFTER THE ACCIDENT AT THE CHERNOBYL NUCLEAR POWER PLANT

ABSTRACT. Background. Fallout from Chernobyl accident was primarily to iodine radioisotopes, with Iodine-131 (I-131) being the most predominant. Radioiodines accumulated following the accident could induce pathologic changes in thyrocytes. Internuclear chromatin bridges and "tailed" nuclei - broken bridge fragments - are considered like cytopathological effects of radiation exposure as these abnormalities are formed from dicentric chromosomes, which are established markers of radiation exposure. **Objective.** To test the possibility that internuclear bridges and tailed nuclei are cytological markers of radiation exposure of the thyroid. **Methods.** We investigated thyrocyte nuclear abnormalities in cytological samples from fine-needle aspiration biopsy in papillary thyroid cancer patients exposed to radioiodine after Chernobyl accident (35 subjects from Gomel region, Belarus) and in papillary thyroid cancer of unexposed patients (25 subjects from Leningrad region, Russia). Nuclear abnormalities included internuclear bridges and "tailed" nuclei were examined. **Results.** Cells in papillary thyroid cancer of irradiated patients are characterized by the high frequency of appearance of hole nucleoplasmic bridges as well as broken bridges in comparison with the control group. The average frequency of thyrocytes with bridges in irradiated patients was almost 4 times higher than that in the unexposed group ($4,69 \pm 0,69\%$ vs. $1,10 \pm 0,23\%$, $p < 0.001$). The same contrast was observed in parameter "frequency of thyrocytes with "tailed" nuclei" ($12,40 \pm 1,82\%$ vs $3,68 \pm 0,39\%$, ($p < 0.001$)). **Conclusion.** Thyrocytes with internuclear bridges may be considered as markers of radiation effects on the thyroid gland.

© V.Iu.Kravtsov, N.V.Ibragimova, S.N.Nikonovich, E.A.Nadyrov, A.V.Rozhko, 2015

✉ natibr@yandex.ru

Citation:

Kravtsov VIu, Ibragimova NV, Nikonovich SN, Nadyrov EA, Rozhko AV. [Internuclear chromosome bridges in thyrocytes of papillary thyroid cancer in patients, subjected to radioactive iodine isotopes during first months after the accident at the Chernobyl nuclear power plant]. *Morphologia*. 2015;9(4):37-42. Russian.




Fig. 1. Internuclear chromosome bridges in thyrocytes. Cytological preparation of papillary thyroid cancer fine-needle biopsy. Romanowsky azure II-eosin staining. $\times 1000$.




Fig. 2. Distribution of patients from Gomel and Leningrad regions by the parameter 'frequency of thyrocytes with internuclear bridges'.




Fig. 3. "Tailed" nucleus – broken bridge in binuclear thyrocyte Cytological preparation of papillary thyroid cancer fine-needle biopsy. Romanowsky azure II-eosin staining. $\times 1000$.




Fig. 4. Distribution of patients from Gomel and Leningrad regions by the parameter 'frequency of thyrocytes with "tailed" nuclei'.

References

1. Gisselsson D, Pettersson L, Höglund M, Heidenblad M, Gorunova L, Wiegant J, Mertens F, Dal Cin P, Mitelman F, Mandahl N. Chromosomal breakage-fusion-bridge events cause genetic intratumor heterogeneity. *Proc Natl Acad Sci U S A*. 2000 May 9;97(10):5357-62.
2. Gisselsson D, Björk J, Höglund M, Mertens F, Dal Cin P, Akerman M, Mandahl N. Abnormal nuclear shape in solid tumors reflects mitotic instability. *Am J Pathol*. 2001 Jan;158(1):199-206.
3. Prokofieva-Belgovskaya AA. [Radiation-induced damage of chromosomes during the early development of Salmon]. *Tsitologiya*. 1961;3(4): 437-45. Russian.
4. Kravtsov Vlu, Proshin SN, Iakovlev AF, Kaminskaia EV, Vakhtin IuB. [Bridges and multipolar mitoses in populations of rat rhabdomyosarcoma RA-23 cells]. *Biull Eksp Biol Med*. 1997 May;123(5):569-72. Russian.
5. Proshin SN, Kravtsov Vlu, Iakovlev AF, Kaminskaia EV, Vakhtin IuB. [Selection in vivo in cell populations of rat RA-23 rhabdomyosarcoma at the trait "frequency of cells with bridges"]. *Genetika*. 1996 Mar;32(3):406-10. Russian.
6. Awa A. Mutation research at ABCC/RERF: cytogenetic studies of atomic bomb exposed populations. *Mutat Res*. 2003 Jan;543(1):1-15.
7. Kravtsov VY, Fedortseva RF, Starkova YV, Yartseva NM, Nikiforov AM. Tailed nuclei and dicentric chromosomes in irradiated subjects. *Appl Radiat Isot*. 2000 May;52(5):1121-7.
8. Anbumani S, Mohankumar MN. Nucleoplasmic bridges and tailed nuclei are signatures of radiation exposure in *Oreochromis mossambicus* using erythrocyte micronucleus cytome assay (EMNCA). *Environ Sci Pollut Res Int*. 2015 Dec;22(23):18425-36. doi: 10.1007/s11356-015-5107-1.
9. Anbumani S, Mohankumar MN. Gamma radiation induced micronuclei and erythrocyte cellular abnormalities in the fish *Catla catla*. *Aquat Toxicol*. 2012 Oct 15;122-123:125-32. doi: 10.1016/j.aquatox.2012.06.001.
10. Nikiforov AM, Fedortseva RF, Monosova EK, Iartseva NM, Kravtsov Vlu. [Nuclei with protrusions--"tailed" nuclei--and radiation cytogenetic markers in a lymphocyte culture after x-ray irradiation]. *Radiats Biol Radioecol*. 2000 May-Jun;40(3):299-304. Russian.
11. Fridman MV, Savva NN, Krasko OV, Zborovskaya AA, Mankovskaya SV, Kurt Werner S, Demidchik YE. Clinical and pathologic features of "sporadic" papillary thyroid carcinoma registered in the years 2005 to 2008 in children and adolescents of Belarus. *Thyroid*. 2012 Oct;22(10):1016-24. doi: 10.1089/thy.2011.0005.
12. Kuzovatov SN, Kravtsov VYu, Vahtin YuB. [Internuclear chromosomal bridges and nuclei with protrusions in cellular populations of rat PA-23 rhabdomyosarcoma]. *Tsitologiya*. 2000;42(11):1097-102. Russian.

13. Abend M, Pfeiffer RM, Ruf C, Hatch M, Bogdanova TI, Tronko MD, Hartmann J, Meineke V, Mabuchi K, Brenner AV. Iodine-131 dose-dependent gene expression: alterations in both normal and tumour thyroid tissues of post-Chernobyl thyroid cancers. *Br J Cancer*. 2013 Oct 15;109(8):2286-94. doi: 10.1038/bjc.2013.574.
14. Abend M, Pfeiffer RM, Ruf C, Hatch M, Bogdanova TI, Tronko MD, Riecke A, Hartmann J, Meineke V, Boukheris H, Sigurdson AJ, Mabuchi K, Brenner AV. Iodine-131 dose dependent gene expression in thyroid cancers and corresponding normal tissues following the Chernobyl accident. *PLoS One*. 2012;7(7):e39103. doi: 10.1371/journal.pone.0039103.
15. Nadyrov E, Rozhko A, Kravtsov V, Mabuchi K, Hatch M, Nakamura N, Nikonovich S, Aleksanin S. Karyopathological traits of thyrocytes and exposure to radioiodines in Belarusian children and adolescents following the accident at the Chernobyl nuclear power plant. *Radiat Environ Biophys*. 2012 May;51(2):187-93. doi: 10.1007/s00411-012-0407-z.
16. Iakovleva IN, Shishkov RV, Poliakov VG, Pankova PA. [Clinico-morphological peculiarities of thyroid cancer among children exposed to the Chernobyl disaster radiation]. *Vopr Onkol*. 2008;54(3):315-20. Russian.
17. Achille M, Boukheris H, Caillou B, Talbot M, de Vathaire F, Sabatier L, Desmaze C, Schlumberger M, Soria JC. Expression of cell cycle biomarkers and telomere length in papillary thyroid carcinoma: a comparative study between radiation-associated and spontaneous cancers. *Am J Clin Oncol*. 2009 Feb;32(1):1-8. doi: 10.1097/COC.0b013e3181783336.
18. Ito M, Bogdanova T, Zurnadzhy L, Saenko V, Rogounovitch T, Mitsutake N, Kondo H, Maeda S, Nakashima M, Tronko M, Yamashita S. Morphological difference in adult thyroid papillary carcinoma between Japan and Ukraine. *Endocr J*. 2014;61(12):1221-8. doi: 10.1507/endocrj.EJ14-0239.
19. Selmansberger M, Feuchtinger A, Zurnadzhy L, Michna A, Kaiser JC, Abend M, Brenner A, Bogdanova T, Walch A, Unger K, Zitzelsberger H, Hess J CLIP2 as radiation biomarker in papillary thyroid carcinoma. *Oncogene*. 2015 Jul 23;34(30):3917-25. doi: 10.1038/onc.2014.311.
20. Takeichi N, Hoshi M, Iida S, Tanaka K, Harada Y, Zhumadilov Z, Chaizhunusova N, Apsalikov KN, Noso Y, Inaba T, Tanaka K, Endo S. Nuclear abnormalities in aspirated thyroid cells and chromosome aberrations in lymphocytes of residents near the Semipalatinsk nuclear test site. *J Radiat Res*. 2006 Feb;47 Suppl A:A171-7.