

A.V.Dniprovskа¹
V.A.Kolesnichenko²
V.O.Fischenko¹

¹ National Pirogov Memorial Medical University, Vinnytsya

² Sytenko Institute of Spine and joint Pathology, Academy of Medical Science, Ukraine, Kharkiv

Key words:

Scheuermann's disease, classic form, atypical form, thoracic kyphosis, lumbar lordosis, correlation analysis.

Received: 24.04.2016
Accepted: 18.05.2016

UDC 616.711-007.54-07:611.08(60.6)

RELATIONSHIP BETWEEN THE PARAMETERS OF SPINE SAGITTAL CONTOUR AND INDICATORS OF FUNCTIONAL STATE OF THE MUSCULOSKELETAL SYSTEM IN PATIENTS WITH SCHEUERMANN'S DISEASE

ABSTRACT. Background. Patients with both forms of Scheuermann's disease are characterized by the spine deformation and low back pain, hypertonus of the extensor muscles with limited mobility of spine and hip, impairment of life quality. However, the relationship of clinical and radiographic parameters studied enough. **Objective.** To establish the correlation of the spine sagittal contour to parameters of the musculoskeletal system functional state in patients with different variants of Scheuermann's disease. **Methods.** Clinical with the definition of the spine mobility (test "fingers - floor"), the thoracic and lumbar spine flexibility (the Schober method), the hip mobility, radiometric with determination of the thoracic kyphosis and lumbar lordosis magnitude (the Cobb method), electromyography with the determination of the amplitude and frequency of biopotentials of right and left half of the m. erector spinae lumbar part, statistics with correlation analysis. **Results.** Different correlations between the studied parameters in classical and atypical forms of Scheuermann's disease were found. **Conclusion.** In patients with the classic form of Scheuermann's disease value of kyphotic spinal deformity directly influence on the spine mobility and the lumbar spine flexibility, disability index, the level of anxiety and pain symptoms and one indirectly affects the pain intensity. In atypical form of Scheuermann's disease value of lumbar lordosis was associated with the pain intensity, and the magnitude of the thoracic kyphosis influenced on a level of anxiety and pain symptoms mainly. These parameters limited spine mobility and range of motion of the hip.

© A.V.Dniprovskа, V.A.Kolesnichenko, V.O.Fischenko, 2016
✉ veakol@rambler.ru

Citation:

Dniprovskа AV, Kolesnichenko VA, Fischenko VO. [Relationship between the parameters of spine sagittal contour and indicators of functional state of the musculoskeletal system in patients with Scheuermann's disease]. *Morphologia*. 2016;10(2):13-7. Ukrainian.




Fig. 1. Regression line of the thoracic kyphosis value on the level of anxiety due to pain expectation (Pain and Anxiety Symptoms Scale) (a) and on the results of "fingers-floor" test in classical form of Scheuermann's disease – approximated curves.




Fig. 2. . Regression line of the pain intensity (Visual Analogue Scale) on the results of "fingers-floor" test and lumbar lordosis value in atypical form of Scheuermann's disease – approximated curves.

References:

1. Weiss HR. Brace treatment can improve thoracic kyphosis during growth: A case report. Open Access J Scie Techn AgiAI Publ. House. 2013;1:6p. doi: 10.11131/2013/100006.

2. Sorenson KM. Scheuermann's Juvenile Kyphosis: Clinical Appearances, Radiography, Etiology and Prognosis. Copenhagen: Munksgaard; 1964, 237 p.
3. Murray PM, Weinstein SL, Spratt KF. The natural history and long-term follow-up of Scheuermann's kyphosis. *J Bone Joint Surg.* 1993; 75-A(2): 236-247.
4. Ristolainen L, Kettunen JA, Heliövaara M, Kujala UM, Heinonen A, Schlenzka D. Untreated Scheuermann's disease: a 37-year follow-up study. *Eur Spine J.* 2012; 21(5):819-824. doi: 10.1007/s00586-011-2075-0.
5. Lucas-García FJ, Vicent-Carsí V, Sánchez-González M. Atypical lumbar Schuermann's disease: a presentation of 6 cases. *Rev Esp Cir Ortop Traumatol.* 2013; 57(2):135-139. doi: 10.1016/j.recot.2012.11.001.
6. Tsirikos AI, Jain AK. Scheuermann's kyphosis: current controversies. *J Bone Joint Surg. (Br).* 2011; 93(7):857-864.
7. Marks VO. [Orthopedic Diagnostics]. Minsk: Science & Technology; 1985, 506 p. Russian.
8. Cobb JR. Outline for the study of scoliosis. *Instruct Course Lectures the Am Acad of Orthop Surgery.* 1948; 5:261-275.
9. Kapandzhy AI. [Lower Limb. Physiologic Anatomy]. Moscow: EKSMO; 2010, 352 p. Russian.